

PAPER • OPEN ACCESS

## Structural resilience of spatial networks with inter-links behaving as an external field

To cite this article: Jingfang Fan *et al* 2018 *New J. Phys.* **20** 093003

View the [article online](#) for updates and enhancements.



**IOP** | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.



## PAPER

# Structural resilience of spatial networks with inter-links behaving as an external field

## OPEN ACCESS

RECEIVED  
3 June 2018REVISED  
5 August 2018ACCEPTED FOR PUBLICATION  
24 August 2018PUBLISHED  
5 September 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.


Jingfang Fan<sup>1,7</sup> , Gaogao Dong<sup>2,3,4</sup>, Louis M Shekhtman<sup>1</sup> , Dong Zhou<sup>1</sup>, Jun Meng<sup>1,7</sup>, Xiaosong Chen<sup>5,6</sup> and Shlomo Havlin<sup>1</sup><sup>1</sup> Department of Physics, Bar Ilan University, Ramat Gan 52900, Israel<sup>2</sup> Institute of applied system analysis, Faculty of Science, Jiangsu University, Zhenjiang, 212013 Jiangsu, People's Republic of China<sup>3</sup> Energy Development and Environmental Protection Strategy Research Center, Faculty of Science, Jiangsu University, Zhenjiang, 212013 Jiangsu, People's Republic of China<sup>4</sup> Center for Polymer Studies and Department of Physics, Boston University, Boston, MA 02215, United States of America<sup>5</sup> CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, People's Republic of China<sup>6</sup> School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China<sup>7</sup> Author to whom any correspondence should be addressed.E-mail: [j.fang.fan@gmail.com](mailto:j.fang.fan@gmail.com) and [jun.meng.phy@gmail.com](mailto:jun.meng.phy@gmail.com)**Keywords:** community structure, percolation, external field, resilience, spatial networksSupplementary material for this article is available [online](#)

## Abstract

Many real systems such as, roads, shipping routes, and infrastructure systems can be modeled based on spatially embedded networks. The inter-links between two distant spatial networks, such as those formed by transcontinental airline flights, play a crucial role in optimizing communication and transportation over such long distances. Still, little is known about how inter-links affect the structural resilience of such systems. Here, we develop a framework to study the structural resilience of interlinked spatially embedded networks based on percolation theory. We find that the inter-links can be regarded as an external field near the percolation phase transition, analogous to a magnetic field in a ferromagnetic–paramagnetic spin system. By defining the analogous critical exponents  $\delta$  and  $\gamma$ , we find that their values for various inter-links structures follow Widom's scaling relations. Furthermore, we study the optimal robustness of our model and compare it with the analysis of real-world networks. The framework presented here not only facilitates the understanding of phase transitions with external fields in complex networks but also provides insight into optimizing real-world infrastructure networks.

## 1. Introduction

Robustness is of crucial importance in many complex systems and plays an important role in mitigating damage [1]. It has been studied widely in both single networks [2–4], interdependent networks [5–10] and multiplex networks [11, 12]. Percolation theory has demonstrated its great potential as a versatile tool for understanding system structural resilience based on both dynamical and structural properties [13, 14], and has been applied to many real systems [15–17]. Recently, a theoretical framework has been developed to study the structural resilience of communities formed of either Erdős–Rényi (ER) and scale-free networks that have inter-links between them using percolation theory [18]. It has been found that the inter-links affect the percolation phase transition in a manner similar to an external field in a ferromagnetic–paramagnetic spin system. However, many real systems, such as, transportation networks [19, 20], infrastructure networks [21] and others, are spatially embedded and the influence of this feature has not been considered. Here we study how the inter-links (e.g. air flights) between two spatial networks (e.g., countries) affect the overall structural resilience. Furthermore, we will search for an optimal structure (or most robust point) of our model and consider it in a real transportation




system. We will do so by developing a framework to study the structural resilience of spatial networks with interlinks and by analyzing possible optimal structures for our model/s and in real transport systems.

The structure of our paper is as follows: in the next section, we describe and introduce the model. In section 3, the results are presented and discussed. Finally, in section 4 a short summary and outlook are provided.

## 2. Model

Our model is motivated by many real-world networks where nodes and links are spatially embedded within the same region (module), but only some nodes have connections to other regions (modules). We denote the links in the same module as *intra*links and the links between different modules as *inter*links. Figure 1(a) demonstrates the topological structure of the global transportation network including railway roads and airline routes [22]. We demonstrate in the figure that the airports are connected via interlinks and can be regarded as *interconnected nodes*. We show here that the interlinks behave, regarding breakdown of the network, in a manner analogous to an external field from physics near magnetic–paramagnetic phase transition [23, 24]. To study this effect, for simplicity and without loss of generalization, we carried out extensive simulations on a network of two modules each with the same number of nodes,  $N_l = L \times L$ , where  $L$  is the linear size of the lattice, representing the spatial networks. Within each module the nodes are only connected with their neighbors in space as defined by a two-dimensional square lattice. Between different modules, we randomly select a fraction  $r$  of nodes to be interconnected nodes, e.g. airports, and randomly assign  $M_{\text{inter}}$  interlinks among nodes in the two modules. A network generated from our model is shown in figure 1(b). Our model is realistic and can represent coupled transport systems, i.e. the nodes in the same lattice module are localized railroad or road networks within the same region while the interlinks represent interregional airline routes.

To quantify the structural resilience of our model, we carried out extensive numerical simulations of the size of the giant connected component  $S(p, r)$  after a fraction of  $1 - p$  nodes are randomly removed. Note that our model is distinct from the case of interdependent networks [5], where the failure of nodes in one network leads to the failure of dependent nodes in other networks. Our model is also different from the interconnected modules model [25], where interconnected nodes are attacked. In our model, the interconnections between different communities are additional connectivity links [26] and randomly chosen nodes are attacked [18]. For a given set of parameters  $[p, r, L]$ , we carried out 10 000 Monte Carlo realizations and took the average of these results to obtain  $S(p, r)$ .



### 3. Results

Similar to our earlier studies [18, 27], we find that the parameter  $r$ , governing the fraction of interconnected nodes, has effects analogous to a magnetic field in a spin system, near criticality. This analogy can be seen through the facts that: (i) the non-zero fraction of interconnected nodes destroys the original phase transition point of the single module; (ii) critical exponents (defined below) of values derived from percolation theory can be used to characterize the effect of external field on  $S(p, r)$ . Figure 2(a) shows our simulation results for the size of the giant component  $S(p, r)$  with  $L = 4096$ ,  $M_{\text{inter}} = 2 \times L \times L$  for various  $r$ . We note that in the limit of  $r = 0$  our model recovers the critical threshold of single square lattices,  $p_c \approx 0.592746$  [28]. We find that  $S(p_c, r) > S(p_c, 0) = 0$  for  $r > 0$ , showing that the interconnected nodes remove the phase transition of the single lattice.

Next, we investigate the scaling relations and critical exponents, with  $S(p, r)$ ,  $p$  and  $r$  serving as our analogy for magnetization (order parameter), temperature, and the external field, respectively [23]. To quantify how the external field,  $r$ , affects the phase transition, we define the critical exponents  $\delta$ , which relates the order parameter at the critical point to the magnitude of the field

$$S(p_c, r) \sim r^{1/\delta}, \quad (1)$$


and  $\gamma$ , which describes the susceptibility near criticality

$$\left( \frac{\partial S(p, r)}{\partial r} \right)_{r \rightarrow 0} \sim |p - p_c|^{-\gamma}, \quad (2)$$

where  $p_c$  is the site percolation threshold for a single two-dimensional square lattice network.

The simulation results for  $\delta$  in our model are shown in figure 2(b). We obtain  $1/\delta = 0.055$  from simulations, which agrees very well with the known exponent value for standard percolation on square lattices  $1/\delta = 5/91$  [13, 14]. The dashed line is the best fit-line for the data with  $R$ -square  $> 0.999$ .

We next investigate the critical exponent,  $\gamma$ , which we claim to be analogous to magnetic susceptibility exponent with the scaling relation given in equation (2). Figure 2(c) presents our results for  $\gamma$ . We obtain  $\gamma = 2.389$  for  $p < p_c$  and  $r = 10^{-4}$ , which agrees again very well with the known value  $\gamma = 43/18$  in




percolation [13, 14]. In figure 2(d) we also plot our results for different  $r$  values:  $r = 10^{-4}$ ,  $10^{-3}$ ,  $10^{-2}$  to highlight the changes in the range of the scaling region. We find that as  $r$  decreases, the scaling region becomes larger, this is expected since for smaller  $r$  the system approaches closer to criticality ( $r = 0$ ). Similar effects in terms of the scaling range are also observed for changing  $M_{\text{inter}}$  with respect to the critical exponent  $1/\delta$  and equation (1), as seen in figure S1 available online at [stacks.iop.org/NJP/20/093003/mmedia](https://stacks.iop.org/NJP/20/093003/mmedia).

We note that for a single 2d square lattice, the scaling exponent  $\beta$ , defined by the relation  $S \sim (p - p_c)^\beta$ , has a value of  $\beta = 5/36$  [13, 14]. The critical exponent  $\beta$  together with  $\delta$  and  $\gamma$  characterize the percolation universality class for our model. Since the various thermodynamic quantities are related, these critical exponents are not independent, but rather can be uniquely defined in terms of only two of them [29]. We find that the scaling hypothesis is also valid for our model and note that our values for these exponents are consistent with the Widom's identity  $\delta - 1 = \gamma/\beta$  [14].

In the following, we test our framework on a real world example involving global transportation networks. We consider two railway networks, one in Europe (EU) and the other in North America (NA). The two railway networks have  $N_{\text{EU}} = 8354$  and  $N_{\text{NA}} = 933$  nodes (stations), as well as  $M_{\text{EU}} = 11128$  and  $M_{\text{NA}} = 1273$  intralinks, respectively. As an example of adding long distance flights, we add  $M_{\text{flight}}$  interconnected links among  $r$  fraction of the nodes (airport hubs). Note that, these nodes are not chosen randomly but based on the real international airports. We used  $M_{\text{flight}} = 1864$ , which is the actual number of direct flights between the two continents. Figure 3 shows our results for the system of the two real networks. We find that, the values of the critical exponents  $\delta$  and  $\gamma$  for the real networks (figures 3(b) and (c)) are consistent with the results obtained from our model. One should note that the percolation threshold  $p_c$  is different in each module when they are separated, since the number of nodes and links is not the same in both modules. To obtain the percolation threshold,  $p_c$  for each real railway network, we analyzed the second largest component,  $S_2(p, 0)$ . The size of the second largest cluster is known to be at a maximum at  $p_c$  [30]. We obtained  $p_c^{\text{EU}} = 0.764$  and  $p_c^{\text{NA}} = 0.758$  by utilizing the peak of  $S_2(p, 0)$  for the EU and NA networks, respectively (see inset of figure 3(a)). For comparison, we also show in figures S2 and S3 the results for the cases where  $r$  nodes are chosen randomly and the links (airlines) between the two spatial sub-networks were assigned randomly. We find that it does not influence the main conclusions: (1) the values of critical exponents are not changed,  $\delta \approx 0.05$  and  $\gamma \approx 2.39$ ; (2) there still exists an optimal amount of interconnected nodes.

To analyze the robustness of our model, we define an effective percolation threshold,  $p_{\text{cut}}$ , by using a small cut-off value of the giant component  $S_{\text{cut}}$  as shown in figure 4(a). The threshold  $p_{\text{cut}}$  is defined as the point where  $S(p, r)$  reaches  $S_{\text{cut}}$ . We assume that when  $S(p, r)$  is very small as  $S_{\text{cut}}$  or below it is not functional. Interestingly, we find an optimal  $r$  in our model. It means that for a certain  $r = r_{\text{opt}}$  the system is most robust i.e.,  $p_{\text{cut}}$  is minimal. Indeed, figure 4(b) shows a specific example with  $S_{\text{cut}} = 0.01$ , where we find the optimal point to be  $r_{\text{opt}} \approx 0.05$ . In our framework, this suggests that if 5% of the cities have interconnected flights the network is most robust to random failures. The origin of this optimization phenomenon is due to the percolation competition between the individual lattice module and the interconnected 'network' composed of  $r$  interconnected nodes/inter-links. When  $r$  is small enough, the behavior of the giant component  $S(p, r)$  is dominated by the single lattice module (see figure 4(a)), and the threshold  $p_{\text{cut}}$  is large and close to  $p_c$  (see figure 4(b), with small  $r$ ); when  $r$  is increasing, the effect of the giant component of a single lattice module becomes weaker, but the effect from the interconnected nodes/inter-links becomes stronger resulting the decreasing of  $p_{\text{cut}}$ ; however, when  $r$  is large, the behavior of the giant component is dominated by the interconnected nodes/inter-links,  $p_{\text{cut}}$  is proportional to  $r$  (see figure 4(b), with large  $r$ ). In particular, our model will become like a random network, when  $r = 1$ . We also




find that, in figure 4(b), there are no significant finite-size effects for our system since the three curves with  $L = 1024, 2048, 4096$  are nearly overlapping. The results on how  $p_{\text{cut}}$  changes with  $S_{\text{cut}}$  and  $r$  are shown in figure 4(c).

Figure 5(a) presents how  $p_{\text{cut}}$  changes with  $S_{\text{cut}}$  and  $r$  for a real network. These results are qualitatively similar to our model results (figure 4(c)). We also observe that there exists an optimal value of  $r$  in the real transportation network. Figure 5(b) shows three specific cases with  $S_{\text{cut}} = 0.01, 0.05, 0.1$ . We find that the optimal point is around  $r_{\text{opt}} \approx 0.1$ . Suggesting that if 10% of cities have intercontinental flights the system is optimally robust against random failures. For comparison, we also show in the figure the fraction of interconnected nodes in the real data:  $r_{\text{EU}} = 0.0055$  and  $r_{\text{NA}} = 0.05$ .

Note that the number of interconnected links,  $M_{\text{inter}}$ , is kept constant when we change  $r$  in our model, i.e.,  $\langle k_{\text{inter}} \rangle$  is proportional to  $1/r$ . We also performed the same analysis to identify how the external field affects the structural resilience, i.e., the critical exponents  $\delta$ ,  $\gamma$  and effective percolation threshold of the spatial and ER networks when  $\langle k_{\text{inter}} \rangle$  is fixed and  $M_{\text{inter}}$  changes, according to  $\langle k_{\text{inter}} \rangle = \langle M_{\text{inter}} \rangle / (rN)$ . The results are presented and discussed in supplemental materials.

#### 4. Summary

In our earlier study [18], we mainly focused on the critical exponents of phase transition in networks with communities, where these communities were not spatially embedded. However, many real systems are spatially embedded. Here, we study the structural resilience of spatial networks (two-dimensional square lattice model) with inter-links and by analyze possible optimal structures for our model/s and in real transport systems. Our model is more realistic and can represent, e.g., coupled transportation systems. In addition, we find that the critical exponents  $\delta$  and  $\gamma$  are constant and do not change with  $\langle k_{\text{inter}} \rangle$  in our spatial networks. However,



**Figure 5.** The effective percolation threshold for a real-world network. (a)  $p_{cut}$  as a function of  $r$  and  $S_{cut}$ . (b)  $p_{cut}$  as a function of  $r$  with  $S_{cut} = 0.01, 0.05, 0.1$ .

different from our model, the value of  $\langle k_{inter} \rangle$  significantly influences the critical exponents in ER networks: only for large  $\langle k_{inter} \rangle$  are equations (1) and (2) satisfied with the mean-field values  $\delta = 2, \gamma = 1$  for the model in [18].

We have developed a framework to study the structural resilience of coupled spatial networks where we show that the inter-links act analogously to an external field in a magnetic–paramagnetic system. Using percolation theory we studied the dynamical evolution of the giant component, and found the scaling relations governing the external field. We defined the critical exponents  $\delta$  and  $\gamma$  using  $S, p$  and  $r$ , which serve as analogs of the total magnetization, temperature and external field, respectively. The values of the critical exponents are universal and relate well with the known values previously obtained for standard percolation on a 2d lattice. Furthermore, we find that our scaling relations obey the Widom’s identity.

We next defined the effective percolation threshold to quantify the robustness of our model. We found that there exists an optimal amount of interconnected nodes, which is also predicted and observed in real-world networks. Our approach provides a perspective on the structural resilience of networks with spatial community structure and gives insight on its response to increasing interlinks in analogy to an external field. Lastly, our model provides a method for optimizing real world interconnected infrastructure networks which could be implemented by practitioners in the field.

## Acknowledgments

We acknowledge the BIU Center for Research in Applied Cryptography and Cyber Security, Italy-Israel project OPERA, the Israel-Italian collaborative project NECST, the Israel Science Foundation, the Major Program of National Natural Science Foundation of China (Grants 71690242, 91546118), ONR, Japan Science Foundation, BSF-NSF, and DTRA (Grant no. HDTRA-1-10-1-0014) for financial support. This work was partially supported by the National Natural Science Foundation of China (Grants 61403171, 71403105, 2015M581738 and 1501100B) and Key Research Program of Frontier Sciences, CAS, Grant No. QYZDJ-SSW-SYS019. JF thanks the fellowship program funded by the Planning and Budgeting Committee of the Council for Higher Education of Israel.

## ORCID iDs

Jingfang Fan  <https://orcid.org/0000-0003-1954-4641>

Louis M Shekhtman  <https://orcid.org/0000-0001-5273-8363>

## References

- [1] Gao J, Barzel B and Barabási A-L 2016 Universal resilience patterns in complex networks *Nature* **530** 307–12
- [2] Cohen R, Erez K, ben Avraham D and Havlin S 2000 Resilience of the Internet to random breakdowns *Phys. Rev. Lett.* **85** 4626–8
- [3] Albert R, Jeong H and Barabási A-L 2000 Error and attack tolerance of complex networks *Nature* **406** 378
- [4] Tanizawa T, Paul G, Cohen R, Havlin S and Stanley H E 2005 Optimization of network robustness to waves of targeted and random attacks *Phys. Rev. E* **71** 047101
- [5] Buldyrev S V, Parshani R, Paul G, Stanley H E and Havlin S 2010 Catastrophic cascade of failures in interdependent networks *Nature* **464** 1025–8
- [6] Leicht E A and D’Souza R M 2009 Percolation on interacting networks arXiv:0907.0894

- [7] Hu Y, Ksherim B, Cohen R and Havlin S 2011 Percolation in interdependent and interconnected networks: abrupt change from second- to first-order transitions *Phys. Rev. E* **84** 066116
- [8] Gao J, Buldyrev S V, Havlin S and Stanley H E 2011 Robustness of a network of networks *Phys. Rev. Lett.* **107** 195701
- [9] Gao J, Buldyrev S V, Stanley H E and Havlin S 2012 Networks formed from interdependent networks *Nat. Phys.* **8** 40–8
- [10] Shekhtman L M, Shai S and Havlin S 2015 Resilience of networks formed of interdependent modular networks *New J. Phys.* **17** 123007
- [11] Hackett A, Cellai D, Gómez S, Arenas A and Gleeson J 2016 Bond percolation on multiplex networks *Phys. Rev. X* **6** 021002
- [12] Solé-Ribalta A, Gómez S and Arenas A 2016 Congestion induced by the structure of multiplex networks *Phys. Rev. Lett.* **116** 108701
- [13] Stauffer D and Aharony A 2003 *Introduction to Percolation Theory* (London: Taylor and Francis)
- [14] Bunde A and Havlin S 2012 *Fractals and Disordered Systems* (New York: Springer)
- [15] Saberi A A 2015 Recent advances in percolation theory and its applications *Phys. Rep.* **578** 1–32
- [16] Li D et al 2015 Percolation transition in dynamical traffic network with evolving critical bottlenecks *Proc. Natl Acad. Sci.* **112** 669–72
- [17] Meng J, Fan J, Ashkenazy Y and Havlin S 2017 Percolation framework to describe El Niño conditions *Chaos* **27** 035807
- [18] Dong G et al 2018 Resilience of networks with community structure behaves as if under an external field *Proc. Natl Acad. Sci.* **115** 6911–5
- [19] Weiss D J et al 2018 A global map of travel time to cities to assess inequalities in accessibility in 2015 *Nature* **553** 333–6
- [20] Strano E et al 2017 The scaling structure of the global road network *R. Soc. Open Sci.* **4** 170590
- [21] Hines P, Blumsack S, Sanchez E C and Barrows C 2010 The topological and electrical structure of power grids 2010 43rd Hawaii Int. Conf. on System Sciences (HICSS) (Piscataway, NJ: IEEE) pp 1–10
- [22] The global railway and airports data are from Natural Earth, <http://naturalearthdata.com/>, The airline routes data are from Openflights, <https://openflights.org/data.html> (Accessed: March 2018)
- [23] Stanley H E 1971 *Phase Transitions and Critical Phenomena* (Oxford: Clarendon)
- [24] Reynolds P, Stanley H and Klein W 1977 Ghost fields, pair connectedness, and scaling: exact results in one-dimensional percolation *J. Phys. A: Math. Gen.* **10** L203
- [25] Shai S et al 2015 Critical tipping point distinguishing two types of transitions in modular network structures *Phys. Rev. E* **92** 062805
- [26] Li D, Qin P, Wang H, Liu C and Jiang Y 2014 Epidemics on interconnected lattices *Europhys. Lett.* **105** 68004
- [27] Shekhtman L M et al 2018 Critical field-exponents for secure message-passing in modular networks *New J. Phys.* **20** 053001
- [28] Newman M E J and Ziff R M 2000 Efficient Monte Carlo algorithm and high-precision results for percolation *Phys. Rev. Lett.* **85** 4104–7
- [29] Domb C 2000 *Phase Transitions and Critical Phenomena* vol 19 (New York: Academic)
- [30] Margolina A, Herrmann H J and Stauffer D 1982 Size of largest and second largest cluster in random percolation *Phys. Lett. A* **93** 73–5